


Impact of Dietary Restrictions on Social Well-being

Student-led Research Conducted by:
Bodhi Chinnaiyan (MI) & Riya Shah (LA)

Introduction


For many people, eating at a restaurant is more than getting a bite. Restaurants are settings for birthdays, family gatherings, friendships, dates, celebrations, and everyday social interactions. For individuals navigating food allergies and other medically necessary dietary restrictions, participation in these experiences can depend on whether restaurants provide sufficient information to support informed dining decisions.

SafeEatsDet collected data from website users in various areas (Michigan, Louisiana, and California) with 35 willing participants who used the website often, illustrating the complexity of this issue at the community level. 57.1% reported personally having a food allergy, while others reported regularly dining with a family member, child, friend, or other individual with an allergy. Among




respondents, restaurant dining was common: 68.6% reported eating at restaurants at least once a month, including 37.1% who reported eating out at least once or twice per week.

Despite the frequency with which respondents encountered restaurant dining, comfort levels varied substantially. 22.9% of respondents reported feeling very uncomfortable eating at restaurants with a food allergy, while an additional 14.3% reported being somewhat uncomfortable. In contrast, 45.7% reported being somewhat or very comfortable. This variation suggests that the restaurant environment does not provide a consistent experience for individuals navigating food allergies.



More importantly, respondents identified a recurring problem: uncertainty. Their concerns included cross-contact, unclear ingredients, hidden allergens in sauces and marinades, inaccurate information from restaurant staff, unclear allergen labeling, limited menu options, and a lack of trust in restaurant practices. In their open-ended responses, participants repeatedly requested clearer ingredient lists, allergen labeling, information about food preparation, and greater transparency regarding cross-contact. Sauces and marinades were among the most frequently mentioned concerns in participants' open-ended responses



These findings are significant because the burden of dietary restriction extends beyond the food itself. When individuals cannot determine whether a meal is safe, dining out requires additional planning, communication, and risk assessment. A restaurant that lacks accessible allergen information may therefore create a barrier to an activity that is fundamentally social.

Dietary Restrictions as a Social and Quality-of-Life Issue

Existing research supports the idea that dietary restrictions can affect quality of life beyond their direct medical consequences. Food allergy has been associated with psychological distress, reduced quality of life, and limitations on everyday activities. Individuals must continually manage the possibility of allergen exposure while navigating environments in which food is present, making dietary restriction an ongoing social consideration rather than an isolated dietary choice.

Research examining individuals with food allergies, celiac disease, and food intolerance has demonstrated that dining outside the home can be particularly challenging. One study found that although eating outside the home was common among individuals with these conditions, participants frequently encountered inadequate knowledge among restaurant staff and concerns regarding the safety of food preparation. Such concerns can make individuals less confident in their ability to eat safely outside their homes.

Therefore, the problem is not simply whether a restaurant offers an allergen-free dish. Individuals must also determine whether the information provided by the restaurant is accurate, whether ingredients are fully disclosed, and whether appropriate precautions are taken during preparation. These uncertainties can transform an ordinary social activity into a situation requiring substantial preparation and vigilance.

Dining Out and Social Participation

The relationship between dietary restrictions and social well-being becomes particularly apparent in research examining dining and social participation. Qualitative research involving individuals with celiac disease has found that maintaining a medically necessary diet can interfere with social activities, travel, and eating outside the home. Participants described difficulties participating in situations where appropriate food was unavailable or where they could not be confident that food had been prepared safely.

Similar findings appear in research involving individuals with multiple food allergies. Participants have described restrictions on spontaneity, stress surrounding food safety, and difficulty participating in social activities centered around food. These findings demonstrate that dietary restrictions can influence not only what individuals consume but also where they go, which activities they participate in, and how comfortable they feel participating in them.

This distinction is important for understanding the SafeEatsDet survey. Only two respondents explicitly reported that an allergy had limited their participation in a social event. However, the survey's open-ended responses revealed substantial concern about restaurant safety and accessibility. Respondents described wanting to know whether restaurants were safe for their allergies, what ingredients were present, how food was prepared, whether cross-contact occurred, and what alternatives were available.

The relatively low percentage reporting direct social restriction should therefore not be interpreted as evidence that restaurant accessibility has little social significance. The question captures whether respondents had already been prevented from participating in a social event, whereas many respondents described barriers that could affect their confidence, restaurant selection, or willingness to dine out. These are related but distinct dimensions of social well-being.

Restaurant Transparency as a Modifiable Barrier

The existing literature suggests that some aspects of the burden associated with dietary restrictions are difficult to eliminate. Individuals with food allergies must continue to avoid allergens, and those with conditions such as celiac disease must maintain medically necessary dietary restrictions. However, the information environment surrounding food is potentially modifiable.

Restaurant communication represents one such area. When menus clearly identify allergens and ingredients, customers can make more informed decisions before ordering. When staff understand allergen protocols and communicate accurately about preparation practices, customers may face less uncertainty. Conversely, unclear menus and inconsistent staff knowledge can increase the cognitive and social burden placed on individuals with dietary restrictions.

The SafeEatsDet survey directly reflects this need. Respondents repeatedly requested ingredient lists, allergen identifiers, clearer menus, information about

preparation methods, cross-contact information, and descriptions of available alternatives. These responses point toward a practical intervention: rather than asking individuals with dietary restrictions to independently navigate an opaque restaurant environment, restaurants and community organizations can make relevant information easier to access.

Implications for SafeEatsDet

The significance of these findings extends beyond documenting that people with food allergies experience difficulties dining out. The survey identifies a specific information gap between individuals who need to make informed dietary decisions and restaurants that possess the information necessary to support those decisions.

SafeEatsDet can address this gap by developing accessible restaurant information focused on the questions that individuals actually need answered before dining: What allergens are present? What ingredients are used? How is the food prepared? What precautions are taken to reduce cross-contact? What alternatives are available? How knowledgeable is restaurant staff about allergen communication?

This approach also shifts the responsibility from simply telling individuals with allergies to "be careful" toward examining the dining environment itself. Individuals will continue to be responsible for managing their medical conditions, but restaurants can reduce unnecessary uncertainty by providing clearer and more consistent information.

The purpose of SafeEatsDet is therefore not to declare restaurants universally "safe" for people with allergies. Such a designation would be medically inappropriate because individual allergies, preparation practices, and risks vary. Instead, the project seeks to increase transparency and informed decision-making by connecting individuals with information that can help them determine whether a restaurant's practices meet their individual needs.


Take-Away

Dietary restrictions can influence social well-being by affecting an individual's ability to participate comfortably and confidently in activities centered around food. Existing research has documented associations between food allergies and other dietary restrictions and reduced quality of life, social limitations, restaurant avoidance, and anxiety surrounding food safety.

The SafeEatsDet survey provides a local perspective on this broader issue. Among the 35 chosen for proven data, about 17% of participants mentioned how allergies limited their dining options and led to worrying.

Respondents repeatedly identified uncertainty surrounding ingredients, cross-contact, labeling, restaurant practices, and staff communication as major concerns. While the survey is too small to establish population-level prevalence, its findings identify a consistent practical problem: people navigating dietary restrictions often need clearer information to participate in restaurant dining with greater confidence.

There are still about 62.8% of individuals who eat out 2 or less times a month, which is less than the American average of 3.



This creates an opportunity for intervention. Improving transparency in restaurant menus, ingredient information, allergen communication, and preparation practices cannot eliminate the medical risks associated with food allergies. It can, however, address an important and modifiable component of the social burden: uncertainty about whether and how a person can participate.

References

SafeEatsDet. <https://www.safeeatsdet.com/>.

Kachru R. Psychosocial issues and quality of life associated with food allergy. *J Food Allergy*. 2020 Sep 1;2(1):95-98. doi: 10.2500/jfa.2020.2.200025. PMID: 39022158; PMCID: PMC11250222.

Kowalczyk A, Moor F. A Meta-Synthesis Exploring Daily Experiences of Adults With Coeliac Disease in Adhering to a Gluten-Free Diet. *J Hum Nutr Diet*. 2025 Apr;38(2):e70043. doi: 10.1111/jhn.70043. PMID: 40197759; PMCID: PMC11977448.

Radke TJ, Brown LG, Hoover ER, Faw BV, Reimann D, Wong MR, Nicholas D, Barkley J, Ripley D. Food Allergy Knowledge and Attitudes of Restaurant Managers and Staff: An EHS-Net Study. *J Food Prot*. 2016 Sep;79(9):1588-1598. doi: 10.4315/0362-028X.JFP-16-085. PMID: 28221943; PMCID: PMC5321626.

Sansweet S, Jindal R, Gupta R. Food allergy issues among consumers: a comprehensive review. *Front Nutr*. 2024 Mar 26;11:1380056. doi: 10.3389/fnut.2024.1380056. PMID: 38595790; PMCID: PMC11002200.

Figuerola-Gómez XA, Oliveras-López MJ, Poyanco-Bugueño MF, Ocaña-Peinado FM, López-García de la Serrana H, Araya Quezada M. Knowledge, attitude, and practices of restaurant and foodservice personnel in food allergy. A systematic review and meta-analysis. *Heliyon*. 2024 Jun 24;10(13):e33431. doi: 10.1016/j.heliyon.2024.e33431. PMID: 39040259; PMCID: PMC11260967.

US Foods. “American Dining Out Habits 2023.” US Foods, <https://www.usfoods.com/tools-tips-and-ideas/articles-and-publications/articles/american-dining-out-habits-2023>.